HAND IN

Answers recorded in question paper

PAGE 1 OF 5 PAGES

QUEEN'S UNIVERSITY FACULTY OF ARTS AND SCIENCE SCHOOL OF COMPUTING

CISC-203* DISCRETE MATHEMATICS FOR COMPUTING SCIENCE

TEST 3 November 2005

Professor Selim G. AKL

Please write your answer to each question only in the box marked Answer. No questions will be answered by the instructor during the exam.
This is a closed-book exam. No computers or calculators are allowed. If you are unsure of what is wanted for a particular question, make a reasonable assumption and write this at the beginning of your answer.
PLEASE NOTE: Proctors are unable to respond to queries about the interpretation of exam questions. Do your best to answer exam questions as written.

NAME: _____

STUDENT NUMBER: _____

FOR INSTRUCTOR'S USE ONLY

Question 1: _____ / 5

Question 2: ____ / 5

Question 3: _____ / 5

Question 4: _____ / 5

TOTAL: _____ / 20

Question 1: [5 marks]

A bowl contains ten red balls and ten blue balls. You select balls at random without looking at them (the selected balls are taken out of the bowl and not replaced).

Answer:

(i) How many balls must you select to be sure of having at least three balls of the same color? Justify your answer.

Hint: Use the pigeonhole principle.

(ii) How many balls must you select to be sure of having at least three blue balls? Justify your answer.

Question 2: [5 marks]

Answer the following questions, justifying your answer in each case. How many strings of six lowercase letters of the 26-letter English alphabet contain

Answer: (i) the letter a, at least once? (ii) the letters a and b, each at least once? (iii) the letters a and b in consecutive positions, such that a immediately precedes b, and all the letters of the string (including a and b) are distinct? (iv) the letters a and b, where a is somewhere to the left of b in the string, and all the letters of the string (including a and b) are distinct?

NAME: _____

PAGE 4 OF 5 PAGES

Question 3: [5 marks]

(a) Let n, r, and k be nonnegative integers, where $r \leq n$ and $k \leq r$. Prove the identity:

 $C(n,r) \times C(r,k) = C(n,k) \times C(n-k,r-k).$

Hint: Expand the two sides of the identity.

Answer:

(b) Let p be a prime and k and integer such that $1 \le k \le p-1$. Show that p divides C(p, k).

Hint: Show that p divides the numerator, but not the denominator.

Answer:

Question 4: [5 marks]

(a) How many different ways are there to choose 12 donuts from the 21 varieties at a donut shop? You can assume that the shop has at least 12 donuts of each variety available. Justify your answer.

Answer:

(b) How many strings of 20 decimal digits are there that contain two 0s, four 1s, three 2s, one 3, two 4s, three 5s, two 7s, and three 9s? Justify your answer.

Answer: